



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


A NEW INFUSORIAN PARASITE IN SAND FLEAS *

MINNIE E. WATSON

Oyster Bay, Long Island, N. Y.

While examining some sand fleas on the shores of Long Island Sound for gregarine parasites, the writer found two hosts infected with an infusorian parasite; one of the hosts being the common small flea, *Orchestia agilis*, the other the larger *Talorchestia longicornis*. The two infected hosts were found in the same habitat and on the same day. The infusoria were found in only these two instances out of about 300 fleas examined for gregarines. The parasites infect the alimentary tract of the host in great numbers, several hundred being present in each flea.

In the vegetative stage, the animals are broadly ovoidal, tapering slightly at one end (Fig. 1). The macronucleus is very large, and ellipsoidal in shape; it is granular and homogeneous and does not



EXPLANATION OF FIGURE

1. *Balantidium orchestium*, vegetative individual. *Og*, oral groove; *oe*, esophagus; *ma*, macronucleus; *mi*, micronucleus; *fv*, food vacuole; *cv*, contractile vacuole.

2, 3, 4. Three stages in binary fission.

5. Conjugation.

stain deeply with Ehrlich's hematoxylin and acetocarmin. The micronucleus is small and deep staining, and lies contiguous to the macronucleus. The apical or subapical oral groove is small and inconspicuous; it leads to a short slender esophagus. Small contractile vacuoles were observed, not more than one being seen in a single individual. Many small food vacuoles were often present. Cilia of uniform length

* Contributions from the Zoological Laboratory of the University of Illinois, under the direction of Henry B. Ward, No. 61. Also from the Biological Laboratory, Cold Spring Harbor.

cover the body, those in the oral groove being slightly the longer. The size of vegetative forms ranges from 300 to 360 microns by 180 to 220 microns.

In individuals which are ready for asexual reproduction, the body becomes elongate and tapers at the ends; the macronucleus becomes longer and constricted centrally, at the same time contracting in volume. It also stains more deeply than does that of the vegetative animal (Figs. 2, 3, and 4). When transverse fission is complete, the two nuclei attain their normal vegetative form and density. Conjugation was noted in one instance.

The parasite may be classified as follows:

1. Order Heterotricha: zone of large cilia leading to mouth.
2. Suborder Polytricha: body covered with an even coat of cilia.
3. Family Bursaridae: peristome broad, body broad and large.
4. Genus *Balantidium*: parasitic, inhabiting the alimentary tract of the host, body ovoidal or ellipsoidal, blunt at ends in vegetative stage, macronucleus ellipsoidal.

5. The parasite is closely allied to *B. coli* Stein and *B. elongatum* Stein; it differs from them in size of the body and in relative size of the nucleus. No previous record of this form has been found.

I wish therefore to designate this species *Balantidium orchestium*.

Urbana, Ill., June, 1915.

REFERENCES CITED

- Doflein, F. 1911. *Lehrbuch der Protozoenkunde*. Jena, 1043 pp.
Minchin, E. A. 1912. *An Introduction to the Study of the Protozoa*. London, 520 pp.